

## Heterosis and heterotic patterns among maize landraces for forage

Lopez Cesar Gabriel<sup>1\*</sup>, Bertoia Luis Maximo<sup>1</sup>, and Burak Ruggero<sup>1</sup>

Received 19 March 2009

Accepted 23 June 2009

**ABSTRACT** - Corn silage is a high-quality forage crop used in many areas of the world. Although vegetative and reproductive components of the plant must be considered, breeding programs in temperate regions are mainly based on the Reid x Lancaster heterotic pattern that has undergone several cycles of improvement for grain yield. Moreover, hybrids selected for forage production are early maturing genotypes not adapted to warm-temperate or subtropical areas. Consequently, exotic germplasm should be considered as a source of materials for breeding programs. Eight landraces were crossed following a diallel mating design. Interpopulation crosses showed high heterosis for ear, stover, and whole plant dry matter yield (EY, SY, and WY, respectively). On average, crosses had higher SY than checks, but lower EY. Considering WY, two interpopulation crosses had higher means than all commercial checks, indicating the potential of the germplasm evaluated. Two composites were selected and different breeding strategies are discussed.

**Key words:** Forage maize, composites, diallel analysis.

### INTRODUCTION

Corn silage is a high-quality forage crop used in many areas of the world, which help dairy and cattle farmers to maintain a relatively constant forage supply during the year. It is used in mass due to its high yield, energy and digestibility.

Even though any maize forage breeders must consider the vegetative and reproductive components of the maize plant (Barriere and Traineau 1986, Dhillon et al. 1990, Argillier et al. 1995), temperate breeding programs largely rely on the use of the Reid x Lancaster heterotic pattern that has undergone several cycles of improvement, primarily for grain yield. Additionally, hybrids developed for superior forage production are early maturing genotypes not adapted to warm-temperate or subtropical areas. Consequently, under these climates, exotic germoplasm should be considered

as a source of material for breeding programs devoted to the development of hybrids with good forage production. According to Crossa et al. (1990), in the Americas there is a tremendous genetic diversity in maize, as a result of thousands of years of evolution under domestication and hybridization, which has not been effectively exploited. Many authors have suggested the usefulness of incorporating exotic germplasm into breeding programs (Eberhart 1971, Hallauer and Miranda 1981, Oyerbides-Garcia et al. 1985, Holley and Goodman 1988, Mungoma and Pollak 1988, Iglesias and Hallauer 1989, Pollak et al. 1991, Michelini and Hallauer 1993, Rodrigues and Chaves 2002, Carena 2005, Soengas et al. 2006).

Thompson (1968) found that a group of exotic and semi-exotic populations yielded on average 28% more digestible dry matter than adapted hybrids, and Stuber

<sup>1</sup> Universidad Nacional de Lomas de Zamora, Facultad de Ciencias Agrarias, C.P.1836, Buenos Aires, Argentina. \*E-mail: cesarglopez63@yahoo.com.ar



(1986) suggested that some semi-exotic materials might be suitable for silage, given their good grain production and great vegetative development. Bosch et al. (1994) showed that some tropical maize populations produced high total digestible dry matter yields in semi-exotic crosses with B73 and MO17 inbred lines. Bertoia (2001) noted that landraces with no history of breeding for grain production generated crosses with good forage potential. Additionally, inbred lines from the North American Corn Belt did not demonstrate potential for enhanced stover yield and quality when compared with inbred lines from Argentine germplasm (Bertoia et al. 2002). According to Vencovsky and Miranda Filho (1972), Miranda Filho (1974), Rodrigues and Chaves (2002), Oliveira et al. (2006), Kutka and Smith (2007), composites are appropriate for use as base populations in breeding programs. Composites are obtained by intercrossing two or more open pollinated varieties with the objective of obtaining a new population with high genetic variability (and a high mean for the traits of interest). Miranda Filho and Chaves (1991) brought the theoretical basis of a procedure for selecting composites based on parameters defined in Gardner and Eberhart (1966) model II for diallel crosses.

The objectives of the present study are i) to select suitable landrace combinations to form composites for forage production adapted to temperate and warm-temperate areas, ii) to propose possible heterotic patterns among them, and iii) to define breeding strategies.

## MATERIAL AND METHODS

Eight maize landraces were evaluated, representing differences in agronomic response, geographic origin, maturity, height, and grain type: ARZM 17-034 (32°13'S - 65°53'W; 906 growing degree days (GDD); 2.64 m, white dent), ARZM 03-056 (29°21'S - 59°59'W; 897 GDD; 2.59 m; white dent), ARZM 01-150 (37°11'S - 62°45'W; 865 GDD; 2.60m; white dent), ARZM 03-054 (29°24'S - 59°41'W; 858 GDD; 2.39 m; white dent), ARZM 16-062 (34°55'S - 67°32'W; 795 GDD; 2.30 m; yellow dent), ARZM 16-042 (33°47'S - 69°03'W; 752 GDD; 2.26 m; orange flint), ARZM 19-006 (37°04'S - 69°09'W; 683 GDD; 2.19 m; orange flint), ARZM 01-088 (38°06'S - 62°14'W; 666 GDD; 2.25 m; orange flint). Seeds were supplied by the Maize Germplasm Bank at INTA Pergamino, Argentina. Landraces were crossed following

a diallel mating design without reciprocals. Crosses were performed in eight isolation blocks. In each isolation block, one population was used as the male and the other seven populations were detasseled and used as females. At least 150 ears per cross were obtained. Landraces per se, the 28 F1 crosses, and four commercial check hybrids (Cargill Semiden 5, Dekalb 4F37, Morgan 369, and Syngenta Pucara, selected for grain production but widely used for forage production in Argentina) were evaluated during two growing seasons (1997-1998; 1998-1999) at Esteban Echeverría (34°83'89" S, 58°84'89" W) and Vicente Casares (35°81'89" S, 58°85'69" W) in the Buenos Aires Province dairy region. Soils are typical Argiudoll (Vicente Casares) and Aquic Argiudoll with silty clay loam and B2t horizon (Esteban Echeverría).

The experimental design was a randomized complete block with three replications within each environment. Experimental units consisted of two 5.20-m rows, spaced 0.70 m apart. Plots were over-planted at 52 seeds per row, then thinned to a density equivalent to 71,500 plants ha<sup>-1</sup> at the three-leaf stage. Each experimental unit was harvested by hand when the kernel milk line in approximately 50% of the plants reached two-thirds of the way down the kernels at the central part of the ear (Hunt et al. 1989). Ear and stover were separated and weighed fresh. A representative sample of each plant component was taken, weighed fresh, and dried with dry forced air, then weighed dry to provide an estimate of dry matter percentage. Stover (SY), ear (EY), and whole plant dry matter yield (WY) were determined. Dried samples were milled to a 1-mm particle size and analyzed with near-infrared reflectance spectroscopy. Near infrared spectra between 1,100 and 2,500 nm at every 2 nm were collected on all milled samples using an NIRS 6500 spectrophotometer (NIRSystem Inc., Silver Spring, MD). In vitro dry matter digestibility of ear (ED) and stover (SD) were predicted by NIRS equations, which were calibrated by the enzymatic method (Gabrielsen 1986). Whole-plant dry matter digestibility (WD) was  $\%WD = \%ED \times 100^{-1} \times HI + \%SD \times 100^{-1} \times (1-HI)$ . Where HI is the forage harvest index ( $EY \times WY^{-1}$ ).

## Diallel cross analysis

Analyses of variance were performed for each variable, using a mixed model where environments and genotype x environment interactions were considered

random effects. Data corresponding to landraces and all possible crosses among them (excluding reciprocals) were analyzed according to Gardner and Eberhart (1966), by the following model:

$$Y_{ij} = U_v + \frac{1}{2}(v_i + v_j) + h_{ij}$$

with  $h_{ij} = h + h_i + h_j + s_{ij}$

where:

$Y_{ij}$  = Mean of the cross between landraces  $i$  and  $j$ ,  
 $U_v$  = mean of all landraces,  $v_i$  = variety effect of landrace  $i$  (difference between the mean of a parent *per se* and the mean of all parents),  $h_{ij}$  = mid-parent heterosis effect,  $h$  = average heterosis (mean of all crosses minus the mean of all landraces),  $h_i$  = variety heterosis effect (heterosis contributed by cultivar  $i$  in those crosses in which it is present, measured as a deviation from the average heterosis effect), and  $s_{ij}$  = specific heterosis effect in the cross between landraces  $i$  and  $j$ .

#### Composite selection

All of the predictions were based on formulas outlined by Miranda Filho and Chaves (1991). The predicted mean of a composite ( $Y_c$ ) of  $k$  components was calculated by:

$$Y_c = U_v + \frac{1}{k} \sum_{i=1}^k v_i + \left(\frac{k-1}{k}\right)h + \frac{2(k-1)}{k} \sum_{i=1}^k h_i + \frac{2}{k} \sum_{i < j=2}^k s_{ij}$$

where:

$k$  = size of the composite (number of landraces), and  $U_v$ ,  $v_i$ ,  $h$ ,  $h_i$ , and  $s_{ij}$  are as defined above. Based on these parameters, the relative contribution of each cultivar to any composite mean can be determined for each composite size ( $k$ ) using the following index (Miranda Filho and Chaves 1991):

$$I_i = \frac{1}{2} v_i + [(k-1)/k] h_i$$

$I_i$  and  $Y_c$  were used to estimate the relative contribution of each cultivar to the mixture mean and to predict the mean of selected composites, respectively. Means across all locations of each year were used to calculate these parameters.

#### RESULTS AND DISCUSSION

According to the combined analysis of variance, genotypes varied significantly ( $P < 0.01$ ) for SY, EY, and WY (Table 1), but not for digestibility traits (SD, ED, and WD, not shown). Variations among checks were observed for SY and EY ( $P < 0.01$ ), but not for WY.

**Table 1.** Analysis of variance for stover (SY), ear (EY), and whole-plant dry matter yield (WY) of eight maize landraces, 28 interpopulation crosses, and four commercial checks

Source of variation	df	SY (Mg ha <sup>-1</sup> )	EY (Mg ha <sup>-1</sup> )	WY (Mg ha <sup>-1</sup> )	
				Mean Squares	Mean Squares
Environments (E)	3	235.2**	307.0**	845.0**	
Replications	8	21.2**	7.9**	48.0**	
Genotypes	35	30.9**	6.4**	43.0**	
Varieties ( $v_j$ )	7	132.8**	14.8**	155.4**	
Heterosis ( $h_{ij}$ )	28	5.4*	4.3*	15.3**	
Average heterosis ( $h$ )	1	51.8*	90.3**	278.9**	
Variety heterosis ( $h_i$ )	7	2.9	0.8	3.3	
Specific heterosis ( $s_{ij}$ )	20	3.9	1.3	6.3	
Checks	3	24.6**	14.4**	6.0	
Checks vs Genotypes	1	983.3**	77.7**	3.0	
Genotype × E	105	3.5**	2.5**	7.0**	
$v_j$ × E	21	4.4**	4.8**	14.0**	
$h_{ij}$ × E	84	3.2**	1.9**	5.8*	
$h$ × E	3	6.3	3.8**	14.9*	
$h_i$ × E	21	3.9**	2.2**	6.9*	
$s_{ij}$ × E	60	2.9**	1.7**	5.0**	
Checks × E	9	4.6**	1.3**	8.0**	
(Checks vs. Genot.) × E	3	11.8**	1.0	20.0**	
Pooled error	312	1.2	0.9	3.0	

\*, \*\*, Significant at 0.05 and 0.01 probability levels for an F test, respectively



The difference between commercial hybrids and experimental genotypes was significant for SY and EY ( $P < 0.01$ ). On average, checks had greater EY but lower SY than the unimproved genotypes (Table 2). No landraces or crosses had EY as high as the best check, Dekalb 4F37 (8,935 kg ha<sup>-1</sup>), but crosses BxG (7,940 kg ha<sup>-1</sup>) and DxG (7,767 kg ha<sup>-1</sup>) did not show significant differences with the second best check (Cargill Semiden 5 with 8,449 kg ha<sup>-1</sup>) and had better EY than Morgan 369 (6,454 kg ha<sup>-1</sup>) (Table 2). SY of crosses AxB (12,883 kg ha<sup>-1</sup>) and AxD (13,229 kg ha<sup>-1</sup>), were significantly greater than the best check, Morgan 369 (11,894 kg ha<sup>-1</sup>). WY ranged from 10,893 kg ha<sup>-1</sup> for landrace H, to 20,329 kg ha<sup>-1</sup> for the cross AxD, which together with cross AxB (19,814 kg ha<sup>-1</sup>) showed significantly higher WY than all checks (Table 2). According to Gardner and Eberhart diallel model II (1966), variety effects ( $v_i$ ), mid-parent heterosis effects ( $h_{ij}$ ), and average heterosis ( $h$ ), were significant for EY, SY, and WY. Specific heterosis ( $s_{ij}$ ) and variety heterosis ( $h_i$ ), did not show significant differences for any trait (Table 1). All effects x environment interactions were significant.

The combined analysis of variance indicates that variety effects accounted for 46% of the entries sum of squares for EY and 86% for SY, while mid-parent heterosis explained 54% and 14% respectively of the entries sum of squares. Landraces B (511 kg ha<sup>-1</sup>), D (754 kg ha<sup>-1</sup>), F (486 kg ha<sup>-1</sup>), and G (508 kg ha<sup>-1</sup>) showed positive and high  $v_i$  values for EY (Table 3). The highest  $v_i$  values for SY were observed in landraces A (1,680 kg ha<sup>-1</sup>), B (2,244 kg ha<sup>-1</sup>), C (727 kg ha<sup>-1</sup>), and D (1,424 kg ha<sup>-1</sup>). High values for varieties effects are indicative of a high frequency of favorable alleles, indicating good potential for the use of these landraces as breeding materials in recurrent selection programs (Crossa et al. 1990).

Crosses showed greater values than parental landraces for all yield traits, indicating significant mid-parent heterosis ( $h_{ij}$ ). With the exception of crosses BxD, CxD and ExF for SY and BxF and CxD for WY, all  $h_{ij}$  effects were significant and relatively high (Table 3), ranging from 6.4 % to 29.2 % for EY, from -8.2 % to 19.4 % for SY, and from 4.3 % to 20.4 % for WY. Besides the high mid-parent heterosis observed in most crosses, high parent heterosis ( $h_{ii}$ , not shown but easily calculated from table 2) must be also taken into account, and only those crosses with high means should be considered. Thus, the best crosses for SY (AxB, and AxD) showed  $h_{ij}$  and  $h_{ii}$  (in parenthesis) values of 10.3% (7.7%) and 17.4% (16.1%) respectively, while for EY,

cross AxD had values of 26.6% (11.1%), BxD 14.7% (12.5%), BxF 20.6% (20.4%), BxG 29.2% (29.1%), BxH 25.5% (16.2%), and FxG 16.6% (16.4%). Considering WY, crosses AxB and AxD showed heterosis values of 15.4% (9.4%) and 20.4% (15.8%), respectively. Several studies reported a high mean,  $h_{ij}$ , and/or  $h_{ii}$  values in interpopulation or line by population crosses (Miranda Filho and Vencovsky 1984, Crossa et al. 1990, Perez-Velazquez 1995, San Vicente et al. 1998, Bertoia 2001, Mickelson et al. 2001, Reif et al. 2003, Soengas et al. 2006).

High means for EY, SY, and WY as well as high  $v_j$  and  $h_{ij}$  effects observed in some landraces and their crosses, make them suitable to be used as a germplasm source in breeding programs. The combination of landraces to form composites could be a good strategy that must be explored.

To evaluate composites among a group of landraces ( $n$ ), the possible number of combinations (assuming equal proportions of each landrace in the composite) is  $N_c = 2^n - (n+1)$  (Vencovsky and Miranda Filho 1972). As an example, with only 10 landraces, 1,013 different combinations can be obtained, making their synthesis and field evaluation prohibitive. Thus, the use of prediction procedures can be very helpful when a large number of entries to evaluate under field conditions is not possible. Miranda Filho and Chaves (1991), proposed a model to select composites. Measuring the relative contribution to the composite means ( $I_i$ ) of each potential landrace to include, they selected only the most promising landraces, thereby reducing the number of possible combinations to test. Furthermore, they calculated the predicted means ( $Y_c$ ) of the chosen combinations as an additional selection criterion.  $Y_c$  is a function of the effects defined in Gardner and Eberhart model II, as can be seen in the equation (1). Those effects are multiplied by weighting coefficients that modify their contribution to the predicted means in a quantity that is a function of the number of landraces ( $k$ ). In this equation (1), for any composite size, the first and third terms are constants, and the fifth term is very small for large  $k$  ( $2/k^2$  is negligible and  $\sum s_{ij}$  tends to 0 when  $k$  tends to  $n$ ). Thus, for each composite size, terms depending on  $v_i$  and  $h_i$  are important in selecting landraces to form composites. The relative contribution of each landrace to the mean of different composites ( $I_i$ ) is defined as a function of both parameters (see equation 2).  $I_i$  and  $Y_c$  were used in this paper to select the most promising composites for forage production.

Heterosis and heterotic patterns among maize landraces for forage

**Table 2.** Mean, and mid-parent heterosis (MPH) for ear, stover, and whole dry matter yield (EY, SY, and WY, respectively), across four environments for landraces, crosses among landraces, and commercial checks. Codes are in letters for landraces and in numbers for commercial checks: A = ARZM 17-034; B = ARZM 03-056; C = ARZM 01-150; D = ARZM 03-054; E = ARZM 16-062; F = ARZM 16-042; G = ARZM 19-006; H = ARZM 01-088; 1 = Morgan 369; 2 = Cargill Semiden 5; 3 = Syngenta Pucara; 4 = Dekalb 4F37

Genotype Code	EY		SY		WY	
	Mean Kg ha <sup>-1</sup>	MPH%	Mean Kg ha <sup>-1</sup>	MPH%	Mean Kg ha <sup>-1</sup>	MPH%
A	4,826	—	11,398	—	16,224	—
B	6,148	—	11,962	—	18,110	—
C	5,574	—	10,445	—	16,019	—
D	6,391	—	11,142	—	17,533	—
E	4,657	—	9,644	—	14,301	—
F	6,123	—	8,820	—	14,943	—
G	6,145	—	8,673	—	14,818	—
H	5,232	—	5,660	—	10,893	—
AxB	6,932	26.3**	12,883	10.3**	19,814	15.4**
AxC	5,532	6.4**	12,634	15.7**	18,166	12.7**
AxD	7,100	26.6**	13,229	17.4**	20,329	20.4**
AxE	5,945	25.4**	11,624	10.5**	17,569	15.1**
AxF	6,490	18.5**	10,453	3.4*	16,943	8.7**
AxG	6,798	23.9**	11,461	14.2**	18,256	17.6**
AxH	6,503	25.6**	9,493	11.3**	15,997	18.0**
BxC	6,679	14.0**	12,580	12.3**	19,259	12.9**
BxD	7,188	14.7**	11,400	-1.3	18,588	4.3*
BxE	6,691	23.9**	11,471	6.2**	18,163	12.1**
BxF	7,400	20.6**	9,534	-8.2**	16,935	2.5
BxG	7,940	29.2**	11,101	7.6**	19,040	15.6**
BxH	7,142	25.5**	9,540	8.3**	16,592	14.4**
CxD	6,598	10.3**	10,658	-1.3	17,256	2.9
CxE	6,646	29.2**	11,454	14.0**	17,901	18.1**
CxF	6,982	19.4**	10,401	8.0**	17,384	12.3**
CxG	6,917	18.0**	10,513	10.0**	17,430	13.0**
CxH	6,531	20.9**	9,617	19.4**	16,148	20.0**
DxE	6,232	12.8**	11,268	8.4**	17,500	9.9**
DxF	6,994	11.8**	11,208	12.3**	18,202	12.1**
DxG	7,767	23.9**	11,522	16.3**	19,289	19.2**
DxH	6,487	11.6**	8,916	6.1**	15,403	8.4**
ExF	6,297	16.8**	9,256	0.3	15,553	6.4*
ExG	6,919	28.1**	9,576	4.6**	16,495	13.3**
ExH	5,564	12.5**	8,723	14.0**	14,288	13.4**
FxG	7,153	16.6**	9,353	6.9**	16,506	10.9**
FxH	6,898	21.5**	7,966	10.0**	14,865	15.1**
GxH	6,511	14.5**	7,678	7.1**	14,189	10.4**
Commercial Checks						
1	6,454	—	11,894	—	18,348	—
2	8,449	—	9,643	—	18,092	—
3	7,498	—	9,297	—	16,794	—
4	8,935	—	8,578	—	17,513	—
LSD (0.05)	752.1	—	893.6	—	1,386.5	—

\*, \*\*, Significantly different from zero at 0.05 and 0.01 probability levels for a t test, respectively



**Table 3.** Estimates of mean of landraces ( $U_v$ ), variety effects ( $v_i$ ), average heterosis ( $h$ ), variety heterosis ( $h_i$ ), and specific heterosis ( $s_{ij}$ ) for ear, stover, and whole plant dry matter yield (EY, SY, and WY, respectively), according to Gardner and Eberhart Model II for a diallel cross among eight landraces (A to H), and predicted means of selected composites

	$v_i$	$h_i$	$s_{ij}$							
			Ear Dry Matter Yield (EY, kg ha <sup>-1</sup> )							
			B	C	D	E	F	G	H	
A	-810.86	95.53	35.68	-649.68	471.97	28.70	-113.27	-103.21	329.81	
B	511.29	213.20		-281.20	-218.95	-3.45	18.65	259.72	189.55	
C	-63.51	-213.90			-94.72	465.82	314.95	-48.61	293.44	
D	754.23	-175.95				-195.86	-119.69	354.52	-197.27	
E	-980.50	-20.56					-105.42	218.23	-408.02	
F	486.18	-67.17						-234.18	238.96	
G	508.45	219.97							-446.47	
H	-405.28	-51.12								
U <sub>v</sub> = 5,637.06 ; h = 1,100.04										
	$v_i$	$h_i$	Stover Dry Matter Yield (SY, kg ha <sup>-1</sup> )							
			B	C	D	E	F	G	H	
			A	1,679.58	480.29	251.16	96.19	633.84	-166.73	-471.01
B	2,243.66	-361.41		601.29	-635.84	240.34	-829.13	231.44	140.74	
C	727.18	303.36			-1,283.66	317.05	130.64	-262.76	401.23	
D	1,423.97	12.17				73.74	880.17	688.78	-357.03	
E	-73.96	-43.46					-267.06	-452.10	254.75	
F	-897.96	-498.22						191.49	364.91	
G	-1,045.16	80.45							-429.00	
H	-4,057.31	26.83								
U <sub>v</sub> = 9,718.08 ; h = 832.78										
	$v_i$	$h_i$	Whole Plant Dry Matter Yield (WY, kg ha <sup>-1</sup> )							
			B	C	D	E	F	G	H	
			A	868.73	575.82	286.84	-553.48	1,105.81	-138.03	-584.28
B	2,754.95	-148.21		320.09	-854.79	236.89	-810.48	491.16	330.29	
C	663.66	89.45			-1,378.38	782.88	445.59	-311.36	694.67	
D	2,178.19	-163.78				-122.11	760.48	1,043.3	-554.30	
E	-1,054.45	-64.02					-372.49	-233.88	-153.26	
F	-411.78	-565.39						-42.69	603.86	
G	-536.71	300.42							-875.47	
H	-4,462.60	-24.29								
U <sub>v</sub> = 15,355.14 ; h = 1,932.82										
Composite	Predicted means									
	EY (kg ha <sup>-1</sup> )	Composite	SY (kg ha <sup>-1</sup> )	Composite	WY (kg ha <sup>-1</sup> )					
B-D	6,729	A-B	12,281	A-B	18,491					
B-F	6,768	A-C	11,778	A-C	17,144					
B-G	7,043	A-D	12,249	A-D	18,604					
D-F	6,566	B-C	11,891	B-C	18,161					
D-G	7,018	B-D	11,476	B-D	18,205					
F-G	6,643	C-D	10,726	C-D	17,016					
B-D-F	6,870	A-B-C	12,222	A-B-C	18,314					
B-D-G	7,164	A-B-D	12,169	A-B-D	18,814					
B-F-G	7,045	A-C-D	11,781	A-C-D	17,920					
D-F-G	6,943	B-C-D	11,425	B-C-D	17,985					
B-D-F-G	7,106	A-B-C-D	11,982	A-B-C-D	18,419					
Average	6,900	Average	11,816	Average	18,098					

$I_i$  of each landrace was estimated for different composite sizes ( $k$ , from 2 to 8), for EY, SY, and WY (Figure 1, 2, and 3). Even though variety heterosis ( $h_i$ ) and specific heterosis ( $s_{ij}$ ) were not significant for any trait, and they were not neglected in the estimates. Similarly, Miranda Filho and Chavez (1991) included  $s_{ij}$  effects in their estimates of  $Y_c$ , although they were non significant.  $I_i$  represents a useful parameter to select the most suitable landraces. For any given trait, positive estimates of  $I_i$  imply that when it is present, the  $i^{th}$  landrace contributes an increasing composite mean. For any  $k$ , Landraces B, D, F, and G had positive  $I_i$  values for EY, and landraces A, B, C, and D for SY. Composite size ( $k$ ) is important because  $I_i$  tends to a general combining ability when  $k$  tends to  $\infty$  (Miranda Filho and Chavez 1991), but depending on the magnitude and sign of  $v_i$  and  $h_i$  effects,  $I_i$  can be positive or negative, or simply increase or decrease, and the relative contribution of each landrace can change along  $k$  values. Landraces A, C, E,

and H showed negative  $I_i$  estimates for EY, and landraces E, F, G, and H for SY. Landraces with negative values of  $I_i$  should be discarded, since they will reduce the mean of any possible composite (with these eight landraces) if included. Two hundred and forty seven composites can be synthesized with these eight landraces, but if those with negative  $I_i$  are discarded, 11 composites for EY and 11 for SY can be formed.

Table 3 shows the predicted mean of the selected composites. Predicted EY means varied from 6,566 kg ha<sup>-1</sup> for composite D-F, to 7,164 kg ha<sup>-1</sup> for composite B-D-G, with an average of 6,900 kg ha<sup>-1</sup> for the 11 composites, similar to the lowest yielding hybrid, Morgan 369 (6,454 kg ha<sup>-1</sup>). The predicted SY ranged from 10,726 kg ha<sup>-1</sup> for C-D, to 12,281 kg ha<sup>-1</sup> for A-B, with an average for the 11 composites of 11,816 kg ha<sup>-1</sup>, similar again to Morgan 369 (11,894 kg ha<sup>-1</sup>) that for SY is the best commercial hybrid. Considering WY, the selected landraces are A, B, C, and D, since they consistently showed positive  $I_i$  estimates. The predicted




Figure 1. Relative contribution ( $I_i$ ) of eight landraces to composite means for EY (kg ha<sup>-1</sup>). Landraces codes are: A= ARZM 17-034; B= ARZM 03-056; C= ARZM 01-150; D= ARZM 03-054; E= ARZM 16-062; F= ARZM 16-042; G= ARZM 19-006; H= ARZM 01-088





Figure 2. Relative contribution ( $I_i$ ) of eight landraces to composite means for SY (kg ha<sup>-1</sup>). Landraces codes are: A= ARZM 17-034; B= ARZM 03-056; C= ARZM 01-150; D= ARZM 03-054; E= ARZM 16-062; F= ARZM 16-042; G= ARZM 19-006; H= ARZM 01-088





**Figure 3.** Relative contribution ( $li$ ) of eight landraces to composite means for WY ( $\text{kg ha}^{-1}$ ). Landraces codes are: A= ARZM 17-034; B= ARZM 03-056; C= ARZM 01-150; D= ARZM 03-054; E= ARZM 16-062; F= ARZM 16-042; G= ARZM 19-006; H= ARZM 01-088

composite WY means varied from  $17,016 \text{ kg ha}^{-1}$  (C-D) to  $18,814 \text{ kg ha}^{-1}$  (A-B-D). An average WY of  $18,098 \text{ kg ha}^{-1}$  show good potential for the selected composites, given that the average actual WY for checks was  $17,687 \text{ kg ha}^{-1}$ , and the best of them (Morgan 369) reached  $18,348 \text{ kg ha}^{-1}$ . Lopez and Mundt (2000), evaluated the ability of the Miranda and Chavez method for predicting

means of wheat mixtures with different number of cultivars (3; 4; and 5), by comparing actual and predicted mixture means for yield (YLD) and diseased leaf area (DLA) under stripe rust infection (*Puccinia striiformis*). They showed significant Spearman's rank correlation coefficients between predicted and actual values varying from 0.78 to 1, under different conditions.

Landraces A, B, C, and D, are good candidates to form a composite, since all have positive  $li$  for WY, SY, and EY, with the exception of A and C for EY. Even considering these negative  $li$  estimates, the results suggest that the best choice is composite A-B-C-D, because the predicted WY ( $18,419 \text{ kg ha}^{-1}$ ) is similar to other good composites of smaller sizes, but including a higher number of landraces will provide higher genetic variability. Additionally, the four populations have similar cycles and grain types, and the predicted mean for EY ( $6,447 \text{ kg ha}^{-1}$ ) is similar to one of the commercial hybrids (Morgan 369,  $6,454 \text{ kg ha}^{-1}$ ). Another composite that can be considered is F-G, formed by two orange flint landraces with similar cycles, positive  $li$  for EY, and combining ability with landrace components of composite A-B-C-D.

Once A-B-C-D is synthesized, the development of recurrent selection schemes followed by the selection of inbred lines can be implemented following two strategies: i) the use of a semi-exotic heterotic patterns among A-B-C-D and inbred lines of well known heterotic patterns such as Reid, Lancaster or other inbred lines characterized by a high potential to increase forage quality (Jung et al. 1998, Argillier et al. 2000), or ii) the exploitation of the exotic heterotic pattern A-B-C-D x F-G, since diallel crosses among them showed high means and heterosis effects for WY, SY, and EY.

#### ACKNOWLEDGEMENTS

Sincere thanks are extended to Dr. Marcelo Ferrer, Germplasm Bank at INTA Pergamino, Argentina, for supplying seeds of the evaluated Landraces.

## Heterose e padrões de heterose entre raças de milho para forragem

**RESUMO** - O milho forrageiro é uma cultura de alta qualidade usada para silagem em muitas áreas do mundo. Embora os componentes vegetativos e reprodutivos da planta devam ser considerados, programas de melhoramento em regiões temperadas, baseiam-se principalmente no padrão heterótico Reid x Lancaster, submetido a vários ciclos de seleção para



melhoramento da produtividade de grãos. Além disso, híbridos selecionados para produção de forragem são genótipos precoces, não adaptados ao calor de zonas temperadas ou subtropicais. Consequentemente, germoplasma exótico deve ser considerado como fonte de genes para programas de melhoramento. Oito raças de milho crioulo foram cruzadas seguindo o modelo diallelo. Cruzamentos interpulacionais apresentaram alta heterose no rendimento de espiga (RE), de palha (RP) e de matéria seca total (RM). Em média, os valores de RP para os cruzamentos foram superiores em relação aos controles; porém inferiores a RE. Com relação a RM, dois cruzamentos interpulacionais tiveram médias superiores a todas as variedades comerciais, indicando o potencial do germoplasma avaliado. Dois compostos foram selecionadas e diferentes estratégias de melhoramento discutidas.

**Palavras-chave:** Milho para silagem, compostos, diallelo.

## REFERENCES

- Argillier O, Méchin V and Barrière Y (2000) Inbred line evaluation and breeding for digestibility related traits in forage maize. *Crop Science* **40**:1596-1600.
- Argillier O, Hébert Y and Barrière Y (1995) Relationships between biomass yield, grain production, lodging susceptibility, and feeding value in silage maize. *Maydica* **40**:125-136.
- Barrière Y and Traineau R (1986) Characterization of silage maize: Patterns of dry matter production, LAI evolution and feeding value in late and early genotypes. In: Dolstra O and Miedema P (eds.) *Proceedings of the 13th Congress on the maize and sorghum section of EUCARPIA*. Wageningen, p. 131-1136.
- Bertoia LM and Peper A (2001) Respuesta forrajera de híbridos de maíz a factores ambientales y de manejo. In: *Actas del VII Congreso Nacional de Maíz*. Pergamino, Argentina.
- Bertoia LM, Burak R and Torrecillas M (2002) Identifying inbred lines capable of improving ear and stover yield and quality of superior silage maize hybrids. *Crop Science* **42**: 365-372.
- Bosch L, Casañas F, Ferret A, Sanchez E and Nuez F (1994). Screening tropical maize populations to obtain semiexotic forage hybrids. *Crop Science* **34**: 1089-1096.
- Carena MJ (2005) Maize commercial hybrids compared to improved population hybrids for grain yield and agronomic performance. *Euphytica* **141**: 201-208.
- Crossa J, Taba S and Welhausen EJ (1990) Heterotic patterns among mexican races of maize. *Crop Science* **30**:1182-1190.
- Dhillon BS, Gurrath PA, Zimmer E, Wermke M, Pollmer WG and Klein D (1990) Analysis of diallel crosses of maize for variation and covariation in agronomic traits at silage and grain harvest. *Maydica* **35**: 297-302.
- Eberhart SA (1971) Regional maize diallels with US and semi-exotic varieties. *Crop Science* **11**: 911-914.
- Gardner CO and Eberhart SA (1966) Analysis and interpretation of the variety cross diallel and related populations. *Biometrics* **22**: 439-452.
- Gabrielsen BC (1986) Evaluation of marketed cellulases for activity and capacity to degrade forage. *Agronomy Journal* **78**: 838-842.
- Hallauer AR and Miranda Filho JB (1981) **Quantitative genetics in maize breeding**. Ames: Iowa State University, 1981. 468p.
- Holley RN and Goodman MM (1988) Yield potential of tropical hybrid maize derivatives. *Crop Science* **28**: 213-218.
- Hunt CW, Kezar W and Vinands R (1989) Yield, chemical composition and ruminal fermentability of corn whole-plant, ear and stover as affected by maturity. *Journal of Production Agriculture* **2**: 357-361.
- Iglesias CA and Hallauer AR (1989) S2 recurrent selection in maize populations with exotic germplasm. *Maydica* **34**: 133-140.
- Jung HG, Mertens DR and Buxton DR (1998) Forage quality variation among maize inbreds: *In vitro* fiber digestion kinetics and prediction with NIRS. *Crop Science* **38**: 205-210.
- Kutka FJ and Smith ME (2007) How many parents give the highest yield in predicted synthetic and composite populations of maize? *Crop Science* **47**: 1905-1913.
- Lopez CG and Mundt CC (2000) Using mixing ability analysis from two way mixtures to predict the performance of cultivars in complex mixtures. *Field Crops Research* **68**: 121-132.
- Michellini LA and Hallauer AR (1993) Evaluation of exotic and adapted maize (*Zea mays* L.) germplasm crosses. *Maydica* **38**: 275-282.
- Mickelson HR, Cordova H, Pixley KV and Bjarnason MS (2001) Heterotic relationships among nine temperate and subtropical maize populations *Crop Science* **41**: 1012-1020.
- Miranda Filho JB (1974) **Cruzamentos dialélicos e síntese de compostos de milho (*Zea mays* L.) com ênfase na produtividade e no porte da planta**. PhD thesis, Escola Superior de Agricultura Luis de Queiroz, Piracicaba, 164p.
- Miranda Filho JB and Vencovsky R (1984) Analysis of diallel crosses among open pollinated varieties of maize (*Zea mays* L.). *Maydica* **29**: 217-234.
- Miranda Filho JB and Chaves LJ (1991) Procedures for selecting composites based on prediction methods. *Theoretical and Applied Genetics* **81**: 265-271.
- Mungoma C and Pollak LM (1988) Heterotic patterns among ten corn belt and exotic maize populations. *Crop Science* **28**: 500-504.

- Oyerbides-García M, Hallauer AR and Cortez-Mendoza H (1985) Evaluation of improved maize populations in Mexico and the U.S. corn belt. **Crop Science** 25: 115-120.
- Oliveira JP, Chaves L J, Duarte JB, Ribeiro KO and Brasil EM (2006) Heterosis for oil content in maize populations and hybrids of high quality protein. **Crop Breeding and Applied Biotechnology** 6: 113-120.
- Perez-Velasquez JC, Ceballos H, Pandey S and Díaz-Amaris C (1995) Analysis of diallel crosses among Colombian landraces and improved populations of maize. **Crop Science** 35: 572-578.
- Pollak L M, Torres-Cardona S and Sotomayor-Rios A (1991) Evaluation of heterotic patterns among caribbean and tropical x temperate maize populations. **Crop Science** 31: 1480-1483.
- Reif JC, Melchinger A E, Xia XC, Warburton ML, Hoisington DA, Vasal SK, Srinivasan G, Bohn M and Frisch M (2003) Genetic distance based on simple sequence repeats and heterosis in tropical maize populations. **Crop Science** 43: 1275-1282.
- Rodrigues MC and Chaves LJ (2002) Heterosis and its components in crosses among high quality protein maize populations. **Crop Breeding and Applied Biotechnology** 2: 281-290.
- San Vicente FM, Bejarano A, Marin C and Crossa J (1998) Analysis of diallel crosses among improved tropical white endosperm maize populations. **Maydica** 43: 147-153.
- Soengas P, Ordás B, Malvar R A, Revilla P and Ordás A (2003) Heterotic patterns among flint maize populations. **Crop Science** 43: 844-849.
- Soengas P, Ordás B, Malvar R A, Revilla P and Ordás A (2006) Combining abilities and heterosis for adaptation in flint maize populations **Crop Science** 46: 2666-2669.
- Stuber CW (1986) Use of exotic sources of germplasm for maize improvement. In: Dolstra O and Miedema P (eds.) **Proceedings of the 13<sup>th</sup> Congress maize sorghum section EUCARPIA**. Wageningen, p. 19-31.
- Thompson DL (1968) Silage yield of exotic corn. **Agronomy Journal** 60: 579-581.
- Vencovsky R and Miranda Filho JB (1972) Determinação do numero de possíveis compostos e pares de compostos. **Relatório Científico do Departamento e Instituto de Genética da Escola Superior de Agricultura Luiz de Queiroz** 6: 120-123.
- Welcker C, Thé C, Andréau B, De Leon C, Parentoni S N, Bernal J, Félicité J, Zonkeng C, Salazar F, Narro L, Charcosset A and Horst W J (2005) Heterosis and combining ability for maize adaptation to tropical acid soils: implications for future breeding strategies. **Crop Science** 45: 2405-2413.